arXiv:1511.07322v1 [astro-ph.IM] 23 Nov 2015

Baltic Astronomy, vol. 24, 000–000, 2015

ASTRONOMICAL SURVEYS AND BIG DATA

Areg M. Mickaelian[1]

[1] *NAS RA V. Ambartsumian Byurakan Astrophysical Observatory (BAO), Byurakan 0213, Aragatzotn Province, Armenia; aregmick@yahoo.com*

Received: 2015 September 1; accepted: 2015 September

Abstract. Recent all-sky and large-area astronomical surveys and their catalogued data over the whole range of electromagnetic spectrum are reviewed, from γ-ray to radio, such as Fermi-GLAST and INTEGRAL in γ-ray, ROSAT, XMM and Chandra in X-ray, GALEX in UV, SDSS and several POSS I and II based catalogues (APM, MAPS, USNO, GSC) in optical range, 2MASS in NIR, WISE and AKARI IRC in MIR, IRAS and AKARI FIS in FIR, NVSS and FIRST in radio and many others, as well as most important surveys giving optical images (DSS I and II, SDSS, etc.), proper motions (Tycho, USNO, Gaia), variability (GCVS, NSVS, ASAS, Catalina, Pan-STARRS) and spectroscopic data (FBS, SBS, Case, HQS, HES, SDSS, CALIFA, GAMA). An overall understanding of the coverage along the whole wavelength range and comparisons between various surveys are given: galaxy redshift surveys, QSO/AGN, radio, Galactic structure, and Dark Energy surveys. Astronomy has entered the Big Data era. Astrophysical Virtual Observatories and Computational Astrophysics play an important role in using and analysis of big data for new discoveries.

Key words: Astronomical Surveys – Astronomical Catalogues – Databases – Archives – Virtual Observatory – Big Data – Laboratory Astrophysics

1. INTRODUCTION

Astronomical surveys are the main source for discovery of astronomical objects and accumulation of observational data for further analysis, interpretation, and achieving scientific results. Recent large multiwavelength (MW) surveys both by ground-based and space telescopes and their catalogues during the last 15 years have accumulated vast amounts of data over the whole range of electromagnetic spectrum from γ-ray to radio. Present astronomical databases and archives contain billions of objects, both Galactic and extragalactic, and the vast amount of data on them allow new studies and discoveries. The Big Data era has come. Astrophysical Virtual Observatories (VO) use available databases and current observing material as a collection of interoperating data archives and software tools to form a research environment in which complex research programs can be conducted. Most of the modern databases give at present VO access to the stored information. This makes possible not only the open access but also a fast analysis and managing of these data. VO is a prototype of Grid technologies that allows distributed data computation, analysis and imaging. Particularly important are data reduction and analysis systems: spectral analysis, spectral energy distribu-

tion (SED) building and fitting, modelling, variability studies, cross-correlations, etc. Numerical or Computational Astrophysics (part of Computer Science, also called Laboratory Astrophysics) has become an indissoluble part of astronomy and most of modern research is being done by means of it. Very often dozens of thousands of sources hide a few very interesting ones that are needed to be discovered by comparison of various physical characteristics. Cross-correlations result in revealing new objects and new samples. The large amount of data requires new approaches to data reduction, management and analysis. Powerful computer technologies are required, including clusters and grids. Large volume astronomical servers have been established to host Big Data and giving high importance to their maintenance, the International Council of Scientific Unions (ICSU) has recently created World Data System (WDS) to unify data coming from different science fields for further possibility of exchange and new science projects.

In our review paper (Mickaelian 2012) "Large Astronomical Surveys, Catalogs and Databases" we summarized astronomical data accumulated during the recent decades and presented astronomical surveys, catalogues, archives, databases and VOs. During these 3 years, many new ground-based and space surveys provided new data and many new releases appeared. Therefore, here we give an update of this information especially emphasizing the importance and going into details of astronomical surveys, as MW photometric ones from gamma-ray to radio, so as proper motion, variability and spectroscopic surveys, including objective prism low-dispersion surveys and digital ones.

2. MULTIWAVELENGTH SURVEYS AND CATALOGUES

Multiwavelength studies significantly changed our views on cosmic bodies and phenomena, giving an overall understanding and posiibility to combine and/or compare data coming from various wavelength ranges. MW astronomy appeared during the last few decades and recent MW surveys (including those obtained with space telescopes) led to catalogues containing billions of objects along the whole electromagnetic spectrum. When combining MW data, one can learn much more due to variety of information related to the same object or area, such as e. g. the Milky Way (Fig 1).

Here we list most important (having homogeneous data for a large number of sources over large area) recent surveys and resulted catalogues providing photometric data along the whole wavelength range:

- ***γ-ray.*** **Fermi-GLAST** 3FGL catalogue: gamma-ray positions and 10 MeV 100 GeV photon counts for 3,033 sources (Acero et al. 2015); **INTEGRAL**: IBIS/ISGRI soft gamma-ray survey catalog of 1,126 sources (Bird et al. 2010); older **CGRO EGRET**: gamma-ray positions and 20 keV 30 GeV photon counts for 1300 sources, including only 271 identified ones (Hartman et al. 1999); **Swift**: survey in deep fields, 9387 sources and **BeppoSAX**: 1082 gamma-ray bursts and some other data are much more accurate, however no all-sky or large-area catalogue is available from these missions/telescopes; typically gamma-ray sources are difficult to identify due to inaccurate positions (several arcmin errors);
- ***X-ray.*** **Röntgensatellit (ROSAT) BSC**: X-ray positions and 0.072.4 keV photon counts and two hardness ratios for 18,806 sources (Voges et al. 1999); **ROSAT FSC**: X-ray positions and 0.07-2.4 keV photon counts and two



Fig. 1. Milky Way image obtained in various wavelengths from γ-ray to radio and showing differences both in individual objects/formations and in the overall shape.

hardness ratios for 105,924 sources (Voges et al. 2000), ROSAT sources are difficult to identify due to inaccurate positions ($\sim 1'$ errors); **INTEGRAL**: hard X-ray all-sky survey catalog of 403 sources; older **EXOSAT** Medium energy (18 keV) Slew Survey Catalog (EXMS): 1210 sources, including 992 identified ones; **ASCA**: 1190 sources and other data are available; much more accurate recent **X-ray Multi-Mirror mission (XMM-Newton)**: 372,728 sources from various surveys and **Chandra**: 380,000 sources from various surveys (no all-sky or large-area coverage);

- ***Ultraviolet (UV).*** **Galaxy Evolution Explorer (GALEX) AIS**: 65,266,291 sources and **GALEX MIS**: 12,597,912 sources, $2''$ rms accurate positions and fluxes in FUV 1528Å and NUV 2271Å bands for some 78 million sources over the whole sky up to 20.8^m for AIS and 22.7^m for MIS (Bianchi et al. 2011); a number of older mission data are available; **EUVE**: 514 sources; **FUSE**: FUV spectra of $\sim$3000 objects and others with much less accuracy and sensitivity; **Hubble Space Telescope (HST)**: 2941Å NUV observations (no large area survey is available);

- ***Optical range.*** **Sloan Digital Sky Survey (SDSS)** Data Release 12 (DR12): 67 mas rms positions, u, g, r, i, z photometry for 932,891,133 objects mainly in Northern extragalactic sky, limited to 22.0^m in u, 22.2^m in g and r, 21.3^m in i and 20.5^m in z (Alam et al. 2015); several POSS I and POSS II based catalogues; **USNO-A2.0**: 526,280,881 objects (Monet et al. 1998), still useful as photometric data differ from USNO-B1.0 and sometimes are better; **Cambridge Automated Plate Measurement (APM)**: $0.6''$ rms positions, b and r photometry for 166,466,987 POSS I objects over 21,000 deg^2, limited to 21.0^m in b and 20.0^m in r (McMahon et al. 2000); **Minnesota Automated Plate Scanner (MAPS)**: $0.5''$ rms positions, O/E photometry for 89,234,404 POSS-I objects over 21,000 deg^2, limited to 21.0^m in O and 20.0^m in E (Cabanela et al. 2003); **USNO-B1.0**: $0.4''$ rms positions, B1, R1, B2, R2, and I photometry; proper motions for all 1,045,913,669 objects present in POSS-I/SERC-J and POSS-II/AAO-SES

over the whole sky, limited to 22.5^m in B and 20.8^m in R; so far the largest among all available catalogues by the number of objects (Monet et al. 2003); and **GSC 2.3.2**: $0.3''$ rms positions, j/V/F/N photometry for 945,592,683 objects present in DSS2, limited to 22.5^m in j and 20.8^m in F (Lasker et al. 2008); **Tycho-2**: 85 mas rms positions and accurate B/V magnitudes for 2,539,913 objects over the whole sky, limited to 15.2^m in B and 16.3^m in V (Hog et al. 2000);

- ***Near-Infrared (NIR).*** **2MASS Point Source Catalog (PSC)**: $0.5''$ rms accurate positions and JHKs photometry for 470,992,970 objects over the whole sky up to 17.1^m in J, 16.4^m in H and 15.3^m in Ks (Cutri et al. 2003); **2MASS Extended Source Catalog (ESC)**: JHK photometry for 1,647,599 objects over the whole sky (Skrutskie et al. 2006); **DEep Near Infrared Survey (DENIS)**: IJKs photometry for 355,220,325 objects over 16,700 deg^2 of the Southern sky up to 18.5^m in I, 16.5^m in J and 14.0^m in Ks (DENIS 2005); deeper NIR observations (no large-area survey) from VLT, UKIRT, and other large IR telescopes;

- ***Mid-Infrared (MIR).*** **Wide-field Infrared Survey Explorer (WISE)**: $0.5''$ rms accurate positions and 3.4 μm (80 μJy sensitivity), 4.6 μm (110 μJy), 12 μm (1 mJy), 22 μm (6 mJy) bands photometry for 563,921,584 sources over the whole sky (Cutri et al. 2012); **AKARI-IRC** Point Source Catalogue: $0.3''$ rms accurate positions and 9 μm (50 mJy sensitivity) and 18 μm (120 mJy sensitivity) fluxes for 870,973 sources over the whole sky (Ishihara et al. 2010); **InfraRed Astronomical Satellite Point Source Catalog (IRAS PSC)** and **Faint Source Catalog (IRAS FSC)**: IR positions and fluxes in 12 μm (0.4 Jy sensitivity) and 25 μm (0.5 Jy) bands for 245,889 sources (IRAS 1988) and 173,044 sources (Moshir et al. 1990), respectively; as well as IRAS joint catalogue has been compiled containing 345,162 sources (Abrahamyan et al. 2015); **IRAS SSSC**: IR positions and fluxes in 12 μm and 25 μm bands for 16,740 extended sources over the whole sky (Helou et al. 1985); **Spitzer Space Telescope (SST)** 7-band data (altogether 4,261,028 sources observed) have accurate positions and much higher sensitivity, however no all-sky or large-area survey is available from this mission;

- ***Far-infrared (FIR).*** **IRAS PSC** and **IRAS FSC**: IR positions and fluxes in 60 μm (0.6 Jy sensitivity) and 100 μm (1.0 Jy) bands for 245,889 sources (IRAS 1988) and 173,044 sources (Moshir et al. 1990), respectively; 345,162 sources in total (Abrahamyan et al. 2015); **IRAS SSSC**: IR positions and fluxes in 60 μm and 100 μm bands for 16,740 extended sources over the whole sky (Helou et al. 1985); **AKARI-FIS** Bright Source Catalogue: $0.8''$ rms accurate positions and fluxes at 65 μm, 90 μm, 140 μm, and 160 μm bands for 427,071 sources over the whole sky with 0.55 Jy sensitivity in 90 μm band (Yamamura et al. 2010);

- ***Sub-mm/mm.*** **Planck**: 30857 GHz (350 μm to 1 cm) 9-band data for the whole sky, study of the CMBR and detection of 33,566 point sources (Planck 2011); **WMAP**: 22-90 GHz fluxes of the whole sky, study of CMBR and detection of 471 point sources (Gold et al. 2011); **SCUBA**: sub-mm instrument on JCMT providing data in 450 μm and 850 μm bands; two large

catalogs have been released: Fundamental Map Object Catalog (JCMTSF), 5061 objects, and Extended Map Object Catalog (JCMTSE), 6118 objects (Di Francesco 2008); **Herschel**: data at 55-672 μm with three instruments in 13 wavelength bands and spectra; **ALMA**: has started providing data and in the nearest future it will produce valuable catalogues in sub-mm/mm wavelengths;

- ***Radio.*** **Green Bank 6 cm (GB6)**: positions and 6 cm (4.83 GHz) fluxes for 75,162 sources sensitive to 18 mJy (Gregory et al. 1996); **NRAO VLA Sky Survey (NVSS)**: positions and 21 cm (1.4 GHz) fluxes for 1,773,484 sources over 33,827 deg^2, sensitive to 2.5 mJy (Condon et al. 1998); **Faint Images of the Radio Sky at Twenty centimeters (FIRST)**: 5″ rms accurate radio positions and 21 cm (1.4 GHz) fluxes for 946,432 sources from 10,000 deg2 area of the Northern extragalactic sky, sensitive to 1 mJy (Helfand et al. 2015); **Sydney University Molonglo Sky Survey (SUMSS)**: 36 cm (843 MHz) fluxes for 211,050 sources over 8000 deg^2 of the Southern sky, sensitive to 1 mJy (Mauch et al. 2012); **Westerbork Northern Sky Survey (WENSS)**: 49 cm (610 MHz) and 92 cm (330 MHz) fluxes for 229,420 sources over 9950 deg^2 of the Northern sky, sensitive to 18 mJy (de Bruyn et al. 1998); **7 Cambridge (7C)**: 198 cm (151 MHz) fluxes for 43,683 sources over 2400 deg^2 of the Northern sky, down to 40 mJy (Hales et al. 2007). Deeper fields are observed on small areas (VLA, ATCA, etc.).

In Table 1 we give a comparative list of multiwavelength all-sky and large-area surveys.

3. OPTICAL IMAGES, PROPER MOTIONS, AND VARIABILITY

Beside the main MW catalogues giving photometric data, there have been a number of astronomical surveys aimed at covering large areas and obtaining optical images (atlases), measuring proper motions or variability. Most important among these surveys and catalogues are:

- ***Optical Images.*** **Digitized Sky Survey (DSS) I**: all-sky digitized images in two bands, blue and red from POSS-I (Palomar Observatory Sky Survey) / SERC-J, 1.67″ sampling (McGlynn et al. 1994); **DSS II**: all-sky digitized images in three bands, blue, red, and IR from POSS-II/AAO-SES, 1″ sampling (Lasker et al. 1996); as mentioned above, most objects are catalogued in USNO B1.0 and GSC 2.3.2; **SDSS DR12**: digital images for 14,555 deg^2 in five bands, u, g, r, i, z, 0.1″ resolution (Alam et al. 2015), this is the most accurate large survey;

- ***Proper Motions.*** **USNO-B1.0**: 0.4″ rms positions and proper motions for all 1,045,913,669 objects over the whole sky (Monet et al. 2003); **GSC 2.3.2**: 0.3″ rms positions for 945,592,683 objects, proper motions were planned, however due to calibration problems they are not listed (Lasker et al. 2008); **Tycho-2**: 85 mas rms positions and proper motions for 2,539,913 objects over the whole sky (Hog et al. 2000); besides, proper motions are accessible based on comparison between POSS I epoch combined measurements from APM/MAPS/USNO-A2.0/USNO-B1.0 and POSS II epoch combined measurements from USNO-B1.0/GSC2.3.2 (Mickaelian & Sinamyan 2010);

Table 1. Main data for the most important all-sky and large-area astronomical surveys providing multiwavelength photometric data. Catalogues are given in the order of increasing wavelengths.

Survey, Catalogue	Years	Spectral range	Sky area (deg^2)	Sensitivity (mag/mJy)	Number of sources	Density (obj/deg^2)
Fermi-GLAST	2008-2014	10MeV-100GeV	All-sky		3,033	0.07
CGRO	1991-1999	20keV-30GeV	All-sky		1,300	0.03
INTEGRAL	2002-2014	15keV-10MeV	All-sky		1,126	0.03
ROSAT BSC	1990-1999	0.07-2.4 keV	All-sky		18,806	0.46
ROSAT FSC	1990-1999	0.07-2.4 keV	All-sky		105,924	2.57
GALEX AIS	2003-2012	1344-2831Å	21,435	20.8^m	65,266,291	3044.85
APM	2000	opt b, r	20,964	21.0^m	166,466,987	7940.61
MAPS	2003	opt O, E	20,964	21.0^m	89,234,404	4256.55
USNO-A2.0	1998	opt B, R	All-sky	21.0^m	526,280,881	12757.40
USNO-B1.0	2003	opt B, R, I	All-sky	22.5^m	1,045,913,669	25353.64
GSC 2.3.2	2008	opt j, V, F, N	All-sky	22.5^m	945,592,683	22921.79
Tycho-2	1989-1993	opt BT, VT	All-sky	16.3^m	2,539,913	61.57
SDSS DR12	2000-2014	opt u, g, r, i, z	14,555	22.2^m	932,891,133	64094.20
DENIS	1996-2001	0.8-2.4 μm	16,700	18.5^m	355,220,325	21270.68
2MASS PSC	1997-2001	1.1-2.4 μm	All-sky	17.1^m	470,992,970	11417.46
2MASS ESC	1997-2001	1.1-2.4 μm	All-sky	17.1^m	1,647,599	39.94
WISE	2009-2013	3-22 μm	All-sky	15.6^m	563,921,584	13669.83
AKARI IRC	2006-2008	7-26 μm	38,778	50 mJy	870,973	22.46
IRAS PSC	1983	8-120 μm	39,603	400 mJy	245,889	6.21
IRAS FSC	1983	8-120 μm	34,090	400 mJy	173,044	5.08
IRAS SSSC	1983	8-120 μm	39,603	400 mJy	16,740	0.42
AKARI FIS	2006-2008	50-180 μm	40,428	550 mJy	427,071	10.56
Planck	2009-2011	0.35-10 mm	All-sky	183 mJy	33,566	0.81
WMAP	2001-2011	3-14 mm	All-sky	500 mJy	471	0.01
GB6	1986-1987	6 cm	20,320	18 mJy	75,162	3.70
NVSS	1998	21 cm	33,827	2.5 mJy	1,773,484	52.43
FIRST	1999-2015	21 cm	10,000	1 mJy	946,432	94.64
SUMSS	2003-2012	36 cm	8,000	1 mJy	211,050	26.38
WENSS	1998	49/92 cm	9,950	18 mJy	229,420	23.06
7C	2007	198 cm	2,388	40 mJy	43,683	18.29

Gaia will measure accurate positions and proper motions for 1 billion stars with an accuracy of about 20 as at 15m, and 200 as at 20m;

- ***Variability.*** Such objects are being obtained by repeated observations and are given in a number of catalogues. **General Catalogue of Variable Stars (GCVS)**: all catalogued 80,671 variables; variability types, typically 0.1m photometric accuracy (Samus et al. 2011); **All Sky Automated Survey (ASAS)**: ASAS Catalogues of Variable Stars (ACVS) containing some 50,000 stars, including some 30,000 new ones (Pojmanski 1998); **Northern Sky Variability Survey (NSVS)**: 14,000,000 objects with 8^m–15.5^m for the area with $\delta > 38^\circ$ (33,326 deg^2); median R ($\sim 0.02^m$ accuracy), measurements scatter, amplitudes, periods, and variability types when possible; 0.04m accuracy (Wozniak et al. 2004a); 8678 catalogued classified red variables from NSVS: Mira type, SR, etc. (Wozniak et al. 2004b); **International Variable Star Index**: the most complete list of stars (altogether 203,438 objects) ever mentioned as variables (www.aavso.org/vsx); besides, variability (as proper motion) studies are possible based on comparison between POSS I epoch combined measurements from APM/MAPS/USNO-A2.0/USNO-B1.0 and POSS II epoch combined measurements from USNO-B1.0/GSC2.3.2 (Mickaelian et al. 2011); **Catalina Sky Survey (CSS)**:

Cataclysmic Variables (Drake et al. 2014a), Periodic Variable Stars (Drake et al. 2014b) and RR Lyrae stars (Abbas et al. 2014); **Panoramic Survey Telescope and Rapid Response System (Pan-STARRS)**: having its major goal to discover and characterize Earth-approaching objects, both asteroids and comets, that might pose a danger to our planet, Pan-STARRS discovers many variable stars, as it performs repeated observations.

Summarizing, we have many times covered the entire sky with optical imaging thus providing data for proper motion and variability measurements. More than 200,000 variable objects have been discovered.

4. SPECTROSCOPIC SURVEYS AND CATALOGUES

Spectroscopic surveys provide important data that can be used for detailed studies of objects. SDSS gives both photometric and spectral data in large area. Most important among these surveys and catalogues are:

- ***Objective Prism Spectroscopy.*** Large amount of spectra have come from objective prism surveys. **First Byurakan Survey (FBS or Markarian Survey)**: 17,000 deg^2 in the Northern sky, 20,000,000 low-dispersion spectra (Markarian et al. 1989); **Second Byurakan Survey (SBS)**: 965 deg^2 in the Northern sky, some 3,000,000 low-dispersion spectra (Stepanian 2005); **Case Low-Dispersion Northern Sky Survey** (Pesch et al. 1995); **Hamburg Quasar Survey (HQS)**: 14,000 deg^2 in the Northern sky (Hagen et al. 1999); **Hamburg-ESO Survey (HES)**: 9,000 deg^2 in the Southern sky (Wisotzki et al. 2000); digitized copies of FBS, HQS, and HES are available online (Mickaelian et al. 2007; Hagen et al. 1999; Wisotzki et al. 2000, respectively); thus, objective prism images for most of the extragalactic sky, 17,000 deg^2 in North and 9,000 deg^2 in South are available;

- ***Medium Dispersion Spectroscopy.*** **2-degree Field and 6-degree Field (2dF/6dF)**: medium dispersion 3700-7900Å spectra for 346,061 galaxies and 49,425 stellar objects, including 23,660 QSOs (galaxy redshift and QSO surveys) (Colless et al. 2001; Croom et al. 2004);

- ***Digital Spectroscopy.*** **SDSS**: 3800-9200ÅÅ R=1800-2200 spectra for 4,355,200 selected objects (mainly galaxy redshift and QSO surveys), including 2,401,952 galaxies, 477,161 QSOs, and 851,968 stars (Alam et al. 2015); **Calar Alto Legacy Integral Field Area (CALIFA)** Survey: is mapping 600 galaxies with imaging spectroscopy (IFS) and produces more than 1 million spectra; **Galaxy and Mass Assembly (GAMA)** spectroscopic survey: is for 300,000 galaxies and also provides millions of spectra.

At present some 5 million objects (compared to 300,000 ones 20 years ago) have spectroscopy giving understanding on their nature and possibility for detailed investigation. The number of QSOs doubles every 5 years (Véron-Cetty & Véron 2010; Pâris et al. 2014). Thousands of Blazars have been identified (Massaro et al. 2015).

A comparative table (Table 2) gives an understanding on various parameters of low-dispersion objective prism surveys and SDSS.

Table 2. Main characteristics of major low-dispersion surveys and SDSS. Most of them are extragalactic surveys so that mainly high galactic latitudes are covered both in the North and South.

Survey	Years	Equipment	Emuls.	D at Hγ	Spectral Range, Å	Covered Area	V_{lim}	Objects of Interest
FBS	1965-1980	102/132cm Schmidt, 1.5° prism	IIa-F	1800	3400-6900	$\|b\|>15°$ $\delta>-15°$ 17,000 deg^2	17.5	UVX (Mkn) galaxies, BSOs, 20 mln spectra
SBS	1978-1991	102/132cm Schmidt, 1.5°/3°/4°	IIIa-J IIIa-F IV-N	1800 900 280	3400-5300 4950-5400 6300-6950	$\|b\|>30°$ $49°<\delta<61°$ 965 deg^2	19	UVX galaxies, QSO/Sy, BCDG, hot stars
Case	1983-1995	61/91cm Schmidt, 1.8° prism	IIIa-J	1350	3400-5300	$\|b\|>30°$ $\delta>30°$	18	BSOs, UVX galaxies CSO/CBS/CG
HQS	1985-1997	80cm Schmidt, 1.7° prism	IIIa-J	1390	3400-5300	$\|b\|>20°$ $\delta>0°$ 14,000 deg^2	19	QSOs, HQS/RASS
HES	1990-1996	1m Schmidt, 4° prism	IIIa-J	280	3400-5300	$\|b\|>30°$ $\delta<+2.5°$ 9,000 deg^2	18	QSOs hot stars
SDSS	2000-2020	2.5m R-C Double MOS	CCD	res.: 2.5Å	3800-9200	$\|b\|>30°$ $\delta>0°$ 14,555 deg^2	21	1 bln objects; 4 mln spectra, 450,000 QSOs

5. BYURAKAN SURVEYS AND THEIR DIGITIZED VERSIONS

The **First Byurakan Survey (FBS)** has been carried out by Markarian, Lipovetski and Stepanian in 1965-1980 with BAO 102/132/213 cm (40″/52″/84″) Schmidt telescope with 1.5° prism (Markarian et al. 1989). 2050 Kodak IIAF, IIaF, IIF, and 103aF photographic plates in 1133 fields (4°×4° each, the size being 16 cm × 16 cm) have been taken. FBS covers 17,000 deg^2 of all the Northern sky and part of the Southern sky (δ > -15°) at high galactic latitudes (|b|>15°). In some regions, it even goes down to δ = -19° and |b|=10°. The limiting magnitude on different plates changes in the range of 16.5^m–19.5^m in V, however for the majority it is 17.5^m–18^m. The scale is 96.8″/mm and the dispersion is 1800 Å/mm near Hγ and 2500 Å/mm near Hβ (mean spectral resolution being about 50Å). Low-dispersion spectra cover the range 3400Å-6900Å, and there is a sensitivity gap near 5300Å, dividing the spectra into red and blue parts. It is possible to compare the red and blue parts of the spectrum (easily separating red and blue objects), follow the spectral energy distribution (SED), notice some emission and absorption lines (such as broad Balmer lines, molecular bands, He, N_1+N_2 lines, broad emission lines of QSOs and Seyferts, etc.), thus making up some understanding about the nature of the objects. The FBS is made up of zones (strips), each covering 4° in declination and all right ascensions except the Galactic plane regions. In all there are 28 zones, which are named by their central declination (ex. zone +27° covers $+25°<\delta<+29°$, zone +63° has $+61°<\delta<+65°$, etc). The zones and the neighboring plates in right ascension overlap about 0.1°, as the exact size of a plate is 4.1°×4.1°, thus making the whole area complete. Each FBS plate contains low-dispersion spectra of some 15,000–20,000 objects, and there are some 20,000,000 objects in the whole survey.

In 1988, Mickaelian continued the survey to low galactic latitudes to check if it is possible to work with the low-dispersion spectra in these crowded regions. Two regions of the Milky Way of the zones +39° and +43° have been covered. 28 Kodak IIaF, IIIaF and 103aF plates in 11 and 8 fields having 171 deg^2 and 117

deg^2 surface respectively have been obtained. These plates are especially useful for discovery of white dwarfs and other galactic objects.

Main features of FBS are:

- First systematic objective-prism survey in the history of astronomy,
- The largest objective-prism survey of the Northern sky (17,000 deg^2),
- New method of search for active galaxies,
- Revelation of 1515 UVX galaxies: some 300 AGN and some 1000 HII galaxies,
- Classification of Seyferts into Sy1 and Sy2 types,
- Definition of Starburst (SB) galaxies,
- FBS Blue Stellar Objects (BSOs) and Late-type Stars,
- Optical identification of IRAS galaxies (BIG and BIS objects); discovery of new AGN and ULIRGs.

The **Second Byurakan Survey (SBS)** was carried out by Markarian, Stepanian, Erastova, Lipovetski, Chavushian, and Balayan in 1978-1991 with BAO 102/132/213cm Schmidt telescope with 1.5°, 3° and 4° prisms, in combination with hypersensitized Kodak IIIaJ and IIIaF plates and filters, giving different ranges of spectrum (3500Å-5400Å, 4950Å-5400Åand 6300Å-6950Å, respectively) (Markarian & Stepanian 1983; Stepanian 2005). In total, 550 plates have been obtained in 65 fields, 4°×4° each. SBS covers 965 deg^2 area at high galactic latitudes ($|b|$ >30°) with $07^h43^m<\alpha<17^h15^m$ and $+49°<\delta<+61°$. The limiting magnitude differs in the range of 18^m–20^m in V, and the survey is complete to 17.5^m. 7 fields (covering 112 deg^2) have limiting magnitudes 19.5^m–20^m (so-called Selected or Deep Fields of the SBS), and are of special interest. The dispersion near Hγ for spectra obtained with 1.5°, 3° and 4° prisms is 1800Å/mm, 900Å/mm, and 280Å/mm, respectively (and 1000Å/mm near Hα for 4° prism). SBS covers 3 zones of the FBS, with central declinations +51°, +55° and +59°. In average, each SBS plate contains low-dispersion spectra of some 50,000 objects, and there are some 3,000,000 objects in the whole survey.

The **Digitized First Byurakan Survey (DFBS)** is the digitized version of the famous Markarian Survey. It is the largest spectroscopic database in the world, providing low-dispersion spectra for 20,000,000 objects. DFBS is a joint project of the Byurakan Astrophysical Observatory (BAO), Cornell University (USA) and Universita di Roma "La Sapienza" (Italy). The whole Northern sky and part of the Southern sky at high galactic latitudes have been observed in FBS, altogether more than 17,000 deg^2. It is especially valuable for extragalactic research. 1500 UV-excess galaxies (Markarian galaxies), 1100 blue stellar objects and 1050 late-type stars have been discovered on the basis of FBS, as well as 1600 infrared (IRAS) sources have been optically identified. The DFBS has been created in 2002-2005 as a result of digitization and reduction of some 2000 FBS plates. High-accuracy (1″ rms) astrometric solution has been made for each plate. Dedicated software allows quick access to any field by given position and extraction of the needed spectra, their calibration, classification and study. The DFBS is free for



Fig. 2. DFBS extraction and analysis software bSpec. It finds objects by an input catalogue data and makes mass extraction and storage of all data with necessary measurements.



Fig. 3. DFBS web interface in mode "Get Spectra". Spectra may be explored and/or extracted in single or batch modes (by the given list).

the astronomical community. It occupies 360 GB space (85 DVDs). The DFBS catalogue and database are being maintained in Armenia and will be also available at CDS, Cornell and Rome. DFBS is the largest Armenian astronomical database and one of the largest in the world. Some projects based on the DFBS have been already put forward, including search for new QSOs and other AGN, continuation of the FBS Second Part, identifications of IR and X-ray sources, etc. The DFBS is the basis for the Armenian Virtual Observatory (ArVO), which will unify all available astronomical data in Armenia, including all Byurakan archive and data from Byurakan telescopes. ArVO is part of the International Virtual Observatories Alliance (IVOA). SBS has also been partially digitized and DSBS is being created.

In Figure 2 we give the DFBS extraction and analysis software bSpec and in Figure 3, DFBS web interface in mode "Get Spectra", where one can select spectra from the list and retrieve corresponding data individually or in batch mode, including 1D FITS spectra.

Table 3. Number of catalogued sources at different wavelength ranges giving a comparative understanding about the wavelength coverage of the observed Universe.

Wavelength range	Major surveys/catalogues	Number of catalogued sources
γ-ray	Fermi-GLAST, INTEGRAL, Swift	10,000
X-ray	ROSAT, XMM-Newton, Chandra	1,500,000
UV	GALEX, HST	100,000,000
Optical	SDSS, DSS I, DSS II	1,700,000,000
NIR	2MASS, DENIS	600,000,000
MIR	WISE, AKARI-IRC, SST	600,000,000
FIR	IRAS, AKARI-FIS, SST, Herschel	500,000
sub-mm/mm	Planck, WMAP, SCUBA, Herschel, ALMA	100,000
Radio	NVSS, FIRST, SUMSS	2,000,000

6. ASTRONOMICAL SURVEYS AND BIG DATA

Astronomical surveys give so much information that huge catalogues, dedicated archives and databases are being built to store, maintain and use these Big Data (Mickaelian 2015). At present astronomers deal with the following numbers in various wavelength ranges (Table 3), and these numbers increase exponentially. It is estimated that there are some 400 billion stars in the Milky Way galaxy and some 100 billion galaxies in the Universe, so that we are very far to catalogue all these objects. Even after Gaia space mission we will have much more accurate astrometric and photometric data for the stars but not much more completeness of detections.

Figure 4 shows the distribution of the numbers of discovered astronomical objects by wavelength range; it is obvious that optical studies are well in front of astronomy together with NIR/MIR (due to 2MASS and WISE), so that even the logarithmic scale does not show the small bars corresponding to other wavelength ranges, beside UV, optical, and NIR/MIR. However MW astronomy was born in the recent decades and makes huge steps toward the overall understanding of the Universe with its various manifestations from γ-ray to radio and in the nearest future most of the objects (e.g. in our Galaxy or all galaxies in the Local Universe) will have their counterparts in all wavelengths.

However, establishing correspondence between sources revealed in different wavelengths is a tricky task. Accurate cross-correlations between various MW catalogues are needed to establish genuine counterparts for each object/sourse. Quick cross-matching is being done for almost all catalogues, however; many objects/sources appear to have false associations, as in crowded regions large contamination with other nighboring objects is happening. Very often individual approach should be applied to such associations. Still, a number of cross-correlation software is in use and is being improved.

We give in Figure 5 a comparison of all-sky and large-area astronomical surveys by their wavelength and sky coverage. It is seen that astronomers have covered the whole sky almost in all wavelength ranges. However, Figure 6 shows that not all wavelength have enough deepness, giving a comparison of all-sky/large-area and deep surveys by their sensitivity (limiting magnitude) and number of catalogued objects, which are rather different (only a few other than optical surveys are seen here; 2MASS, DENIS, WISE, GALEX, and IRAS, others having much smaller numbers).

Large astronomical surveys have become one of the most important directions of investigations in our science and they provide the main bulk of information that has been transformed into Big Data and approached astronomy and computer



Fig. 4. Distribution of the numbers of discovered astronomical objects by wavelength range. Numbers in optical, NIR/MIR and UV are so big that even the logarithmic scale does not show the small bars corresponding to γ-ray, X-ray, FIR, sub-mm/mm and radio.



Fig. 5. Comparison of all-sky and large-area surveys by wavelength range and survey area. There are all-sky surveys almost in all wavelength domains.

science posing new problems and inquiring new solutions.

REFERENCES

Abbas M. A., Grebel E. K., Martin N. F., et al. 2014, MNRAS 441, 1230
Abrahamyan, H. V.; Mickaelian, A. M.; Knyazyan, A. V. 2015, A&C 10, 99
Acero, F.; Ackermann, M.; Ajello, M.; et al. 2015, ApJS 218, 23
Alam, S.; Albareti, F. D.; Allende Prieto, C.; et al. 2015, ApJS 219, 12
Bianchi, L.; Herald, J.; Efremova, B.; et al. 2011, Ap&SS 335, 161
Bird, A. J.; Bazzano, A.; Bassani, L.; et al. 2010, ApJS 186, 1
Cabanela, J. E., Humphreys, R. M., Aldering, G., et al.: 2003 PASP 115, 837
Colless, M.; Dalton, G.; Maddox, S.; 2001, MNRAS 328, 1039



Fig. 6. Comparison of all-sky and large-area surveys by limiting magnitude and number of objects. Only a few other than optical surveys are among those having the biggest numbers (hundred millions and billion objects).

Condon, J. J., Cotton, W. D., Greisen, E. W.; et al.: 1998 AJ 115, 1693
Croom, S. M.; Smith, R. J.; Boyle, B. J.; et al. 2004, MNRAS 349, 1397
Cutri, R. M.; Skrutskie, M. F.; van Dyk, S.; et al. 2003, IPAC/California Institute of Technology
Cutri, R. M.; Wright E. L.; Conrow T.; et al. 2012, WISE All-Sky DR, IPAC/Caltech, VizieR Catalog II/311
DENIS consortium 2005, 3rd release, Vizier online data catalogue II/263
de Bruyn G., Miley G., Rengelink R., et al. 1998, WENSS Collab. NFRA/ASTRON and Leiden Obs., Vizier online catalogue VIII/62
Di Francesco, J. 2008, AAS 212, 9603
Drake A. J., Gansicke B. T., Djorgovski S. G., et al. 2014a, MNRAS 441, 1186
Drake A. J., Graham M. J., Djorgovski S. G., et al. 2014b, ApJS 213, 9
Gold, B., Odegard, N., Weiland, J.L., et al. 2011, ApJS 192, 15
Gregory, P. C., Scott, W. K., Douglas, K., Condon, J. J. 1996, ApJS 103, 427
Hagen, H.-J.; Engels, D.; Reimers, D. 1999, A&AS 134, 483
Hales, S. E. G.; Riley, J. M.; Waldram, E. M.; et al. 2007, MNRAS 382, 1639
Hartman, R. C.; Bertsch, D. L.; Bloom, S. D.; et al. 1999, ApJS 123, 79
Helfand, D. J.; White, R. L.; Becker, R. H. 2015, ApJ 801, 26
Helou, G.; Walker, D. W. 1985, IRAS small scale structure catalog, JPL, Pasadena, NASA
Hog, E., Fabricius, C., Makarov, V. V., et al. 2000, A&A, 355, L27
IRAS, 1988, Joint IRAS Scienve Working Group. IRAS PSC, Version 2.0, NASA RP-1190
Ishihara, D.; Onaka, T.; Kataza, H.; et al. 2010, A&A 514, 1
Lasker, B. M., Doggett, J., McLean, B., et al. 1996, ASP Conf. Ser. 101, 88
Lasker, B. M., Lattanzi, M. G., McLean, B. J., et al. 2008, AJ, 136, 735L
Markarian, B. E.; Lipovetsky, V. A.; Stepanian, J. A.; et al. 1989, Comm. SAO 62, 5
Massaro, E.; Maselli, A.; Leto, C.; et al. 2015, Ap&SS 357, 75
Mauch T., Murphy T., Buttery H.J., et al. 2012, MNRAS 342, 1117, 2003; Online

version 2.1r, 16.02.2012
McGlynn, T., White, N. E., Scollick, K., 1994, ASP Conf. Ser., 61, 34
McMahon, R.G., Irwin, M.J., Maddox, S.J. 2000, The APM-North Catalogue, Inst. of Astron., Cambridge, UK
Mickaelian, A. M. 2012, Baltic Astronomy 21, 331
Mickaelian, A. M. 2015, Astronomy Report, in press
Mickaelian, A. M.; Nesci, R.; Rossi, C.; et al. 2007, A&A 464, 1177
Mickaelian A.M., Mikayelyan G.A., Sinamyan P.K. 2011, MNRAS 415, 1061
Mickaelian A., Sinamyan P. 2010, MNRAS 407, 681
Monet, D.; Bird, A.; Canzian, B.; et al. 1998, USNO Flagstaff Station and Universities Space Research Association (USRA), Vizier online catalogue I/252
Monet, D. G.; Levine, S. E.; Canzian, B.; et al. 2003, AJ 125, 984
Moshir, M., Kopan, G., Conrow, T., et al. 1990, IRAS FSC, Version 2.0, NASA
Pâris, I.; Petitjean, P.; Aubourg, É.; et al. 2014, A&A 563, 54
Pesch, P.; Stephenson, C. B.; MacConnell, D. J. 1995, ApJS 98, 41
Planck 2011, Planck Early Release Compact Source Catalogue Planck Collaboration, ESA, 2011, Vizier catalog VIII/88
Pojmanski G. 1998, Acta Astron. 48, 35
Samus, N. N., Durlevich, O. V., Zharova, A. V., et al. 2011, GCVS database, Inst. Astron. and SAI, Moscow
Skrutskie, M. F.; Cutri, R. M.; Stiening, R.; et al. 2006, AJ 131, 1163
Stepanian, J.A. 2005, RMxAA 41, 155
Tsvetkov, M.K., Stavrev, K.Y., Tsvetkova, K.P., et al. 1994, Proc. IAU S161, Kluwer, Dordrecht, p. 359
Véron-Cetty, M.-P.; Véron, P. 2010, A&A 518, 10
Voges, W., Aschenbach, B., Boller, T., et al. 1999, A&A, 349, 389
Voges, W., Aschenbach, B., Boller, T., et al. 2000, MPE Garching
Wisotzki, L.; Christlieb, N.; Bade, N.; et al. 2000, A&A 358, 77
Wozniak, P. R., Vestrand, W. T., Akerlof, C. W., et al. 2004, AJ 127, 2436
Yamamura, I.; Makiuti, S.; Ikeda, N.; et al. 2010, AKARI/FIS All-Sky Survey Point Source Catalogues